



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES—UNTABULATED.

ARGENTINA—*Buenos Aires*.—Month of September, 1909. Estimated population, 1,214,646. Total number of deaths, 1,629, including diphtheria 4, measles 13, scarlet fever 3, typhoid fever 6, whooping cough 2, and 174 from tuberculosis.

CANADA—*Quebec—Sherbrooke*.—Month of November, 1909. Estimated population, 15,300. Total number of deaths, 16, including 2 from tuberculosis.

FRANCE—*St. Etienne*.—Two weeks ended November 15, 1909. Estimated population, 150,000. Total number of deaths, 122, including scarlet fever 1, typhoid fever 1, and 22 from tuberculosis.

GREAT BRITAIN AND IRELAND.—Week ended November 13, 1909.

England and Wales.—The deaths registered in 76 towns having an aggregate estimated population of 16,445,281, correspond to an annual rate of 13.8 per 1,000.

Ireland.—The deaths registered in 21 principal town districts having an aggregate estimated population of 1,142,308, correspond to an annual rate of 19.2 per 1,000. The lowest rate was recorded at Drogheda, viz, 4.1, and the highest at Portadown, viz, 31 per 1,000.

Scotland.—The deaths registered in 8 principal towns having an aggregate estimated population of 1,865,571 show a total of 566, including diphtheria 7, measles 22, scarlet fever 10, typhoid fever 4, typhus fever 1, and 5 from whooping cough.

ITALY—*Genoa*.—Two weeks ended October 31, 1909. Estimated population, 279,163. Total number of deaths, 165, including diphtheria 1, smallpox 1, typhoid fever 1, and 29 from tuberculosis. Two weeks ended November 15, 1909. Total number of deaths, 177, including diphtheria 3, and 23 from tuberculosis.

PORTO RICO.—Month of November, 1909. Estimated population, 1,053,963. Total number of deaths, 184, including diphtheria 3, measles 2, smallpox 10, typhoid 17, and 151 from tuberculosis.

SAMOA—*Tutuila*.—Month of September, 1909. Estimated population, 6,200. Total number of deaths, 17, including 1 from tuberculosis. Month of October, 1909. Total number of deaths, 20, including 1 from tuberculosis.

SPAIN—*Almeria*.—Month of October, 1909. Estimated population, 50,910. Total number of deaths not reported. Five deaths from typhoid fever, 4 from measles, 3 from scarlet fever, and 3 from smallpox reported.

Seville.—Month of October, 1909. Estimated population, 148,315. Total number of deaths, 367, including diphtheria 1, scarlet fever 1, typhoid fever 1, and 77 from tuberculosis.